

## Sample Paper-II (Unsolved)

Subject : Chemistry

Tin	e: 3 Hrs.	Class: IX	M.M.: 70
Ger	eral Instructio	ns:	
(i) (ii) (iii) (iv) (v) (vi) (vii)	All questions at Marks for ques Question no. 1- Questions no. 5 Question no. 19 Question no. 26 Answer should	re compulsory. tions are indicated against each. -8 are very short answer questions can 9-18 are short answer questions carryi 9-27 are also short answer question ca 8-30 are long answer questions carryi be brief and to the point.	ing 2 marks each. nrrying 3 marks each. ng 5 marks each.
1.		ect set of four quantum numbers for t assium $(Z = 19)$ .	he valence
2.	_	C name and symbol for the element	1
3.	What property van der Walls	of molecules of real gases is indicate constant 'a'?	ted by
4.	/ // //	work was done on a system, it lost 12 the value of internal energy change	
5.	Write K <sub>p</sub> is term	ns of K <sub>c</sub> for the following chemical rea	ction: 1
		$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$	
6. 7.		ple of disproportionation reaction.  ope of hydrogen which contains equa	l number of protons
8.	Write IUPAC n	ame of the following organic compun	d:
	CH <sub>3</sub> -C(Cl	$H_3)_2 - CH = CH_2$	1
9.	with chlorofor	inking water was found to be severel m (CHCl <sub>3</sub> ), supposed to be carcinoge nination was 15ppm (by mass).	-
	a) Express thi	s in percent by mass.	
		the molality of chloroform in the water	er sample.
	[Given molar n	nass of CHCl3 = 118.5g-mol-1	2

## Visit www.ncerthelp.com For All NCERT solutions, CBSE sample papers, Question papers, Notes for Class 6 to 12



- 10. An electron has a speed of  $40 \text{ms}^{-1}$  accurate upto 99.99%. What is the uncertainty in locating its position? [Given  $m_e = 9.11 \times 10^{-3}1 \text{ kg}$ ]
  - 2

11. Give correct reason for the following:

- 2
- a) BF<sub>3</sub> has a zero dipole moment although the B-F bonds are plar.
- b) All carbon to oxygen bonds in CO<sub>3</sub><sup>2</sup>- are equivalent.
- Balance the following redox reaction by ion electron method (in basic medium):

$$MnO_4^-(aq) + \Gamma(aq) \rightarrow MnO_2(s) + I_2(s)$$

13. Complete the following reactions:

2

2

- a) PbS  $(s) + H_2O_2(aq) \rightarrow$
- b)  $Ca(HCO_3)_2 + Ca(OH)_2 \rightarrow$

OR

Account for the following:

- a) Soft water lathers with soap but not hard water.
- b) Temporary hardness of water can be removed by boiling.
- a) Mention the type of hybridisation of each carbon in the compound CH<sub>3</sub> – CN.

- b) Draw the structure of 2-methylpropan-2-ol.
- 15. In sulphur estimation, 0.157g of an organic compound gave 0.4813g 2 of barium sulphate. What is the percentage of sulphur in the compound?
- 16. Write the structures of products (A and B) of the following reactions:
- 2

a) 
$$HC \equiv CH \xrightarrow{Na} A \xrightarrow{CH_3Br} B$$

b) BrH<sub>2</sub> C - CH<sub>2</sub> Br 
$$\xrightarrow{\text{Alcohol}}$$
 A  $\xrightarrow{\text{NaNH}_2}$  B

- 17. Arrange the following in order of property mentioned against each:
- 2
- a) BeCl2, MgCl2, CaCl2, BaCl2 [Increasing ionic character]
- b) Mg(OH)2, Ca(OH)2, Ba(OH)2, Sr(OH)2 [Increasing solubility in water]
- 18. Give reasons for the following:
  - a) The enthalpy of neutralisation is always constant *i.e.*, 57.1 kJ/mol when a strong acid neutralises a strong base.
  - b) Neither q nor W is a state function but q + W is a state function.

## Visit www.ncerthelp.com For All NCERT solutions, CBSE sample papers, Question papers, Notes for Class 6 to 12



- 19. a) State Gay Lussac's law of gaseous volumes.
  - b) What mass of CaCO<sub>3</sub> is required to react completely with 25ml of 0.75M HCl according to equation?

$$CaCO_3(s) + 2HCl(aq) \rightarrow CaCl_2(aq) + CO_2(g) + H_2O(l)$$

[Given atomic masses: Ca + 40u, C = 12u, O = 16u, Cl = 35.5u and H = 1u]

- 20. Account for the following:
  - a) Ionization enthalpy of nitrogen is more than that of oxygen.
  - b) A cation is always smaller then its parent atom.
- 21. Calculate the standard enthalphy of formation of CH<sub>3</sub>OH (*l*) from the following data:

CH<sub>3</sub>OH (*l*) + 
$$\frac{3}{2}$$
 O<sub>2</sub> (*g*)  $\rightarrow$  CO (*g*) + 2H<sub>2</sub>O (*l*) ;  $\triangle$ H<sup>0</sup> = 726 kJ/mol

C (graphite) + 
$$O_2$$
 (g)  $\rightarrow$   $CO_2$  (g) ;  $\Delta_1H^0 = 393 \text{ kJ/mol}^{-1}$ 

$$H_{2}(g) + \frac{1}{2} O_{2}(g) \rightarrow H_{2}O(l)$$
 ;  $\Delta H^{0} = 286 \text{ kJ/mol}$ 

3

- 22. a) State Pauli's exclusion principle.
  - b) Account for the following:
     i) Chromium has configuration 3d<sup>5</sup> 4s<sup>1</sup> and not 3d<sup>4</sup> 4s<sup>2</sup>.
    - ii) Bohr's orbit are called stationary states.
- 23. Explain the following terms:
  - a) Photochemical smog
  - b) Acid rain
  - c) Eutrophication
- a) Write the molecular orbital configuration of O<sub>2</sub><sup>+</sup>. Calculate its bond order and predict its magnetic behaviour.
  - b) What is the state of hybridisation of nitrogen in NH<sub>4</sub><sup>+</sup>ion? 3
- 25. Account for the following:
  - a) Potassium carbonate cannot be prepared by Solvay process.
  - b) Beryllium and magnesium do not give colour to flame.
  - Alkali metals and alkaline earth metals cannot be obtained by chemical reduction methods.

OR

What happens when:

- a) sodium metal is dropped in water?
- b) sodium metal is heated in free supply of air?
- c) sodium peroxide dissolves in water?
- 26. a) What would be the S.I. unit for quantity  $pV^2T^2/n$ ?

3

- b) Calculate the volume occupied by 8.8g of CO<sub>2</sub> at 31.1°C and 1 bar pressure. (R = 0.083 bar LK<sup>-1</sup> mol<sup>-1</sup>).
- 27. a) Arrange the following in order of property mentioned against each: 3
  - i) (CH<sub>3</sub>)<sub>3</sub>C<sup>6</sup>, CH<sub>3</sub>CH<sub>2</sub>CH<sup>+</sup> CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sup>+</sup><sub>3</sub> [Increasing stability order]
  - ii) CH,CH,COOH, (CH,), CHOOH and (CH,), CCOOH

[Increasing acidic strength]

- b) Write the name of isomerism among the following compounds: CH<sub>3</sub>O CH<sub>2</sub>CH<sub>3</sub> and CH<sub>3</sub>CH<sub>2</sub>O CH<sub>3</sub> CH<sub>3</sub>
- 28. a) Write the conjugate acid of HCOO.

5

- b) Calculate the pH of a  $1.0 \times 10^{-8}$  M solution of HCl.
- c) Calculate the solubility of  $A_2X_3$  in pure water, assuming that neither kind of ion reacts with water. The solubility product of  $A_2X_3$  is  $K_{\infty} = 1.1 \times 10^{-23}$ .

OR

- a) Write the conjgate acid of NH<sub>3</sub>.
- b) Assign reason for the following:
  - i) A solution of NH<sub>4</sub>Cl in water shows pH less than 7.
  - ii) In qualitative analysis NH<sub>4</sub>Cl is added before adding NH<sub>4</sub>OH for testing Fe<sup>3+</sup> or Al<sup>3+</sup> ions.
- c) Consider the reaction ₩

$$N_2(g) + 3H_2(g)$$
 2NH<sub>3</sub>+Heat

Indicate the direction in which the equilibrium will shift when:

- i) temperature is increased.
- ii) pressure is increased.

## Visit www.ncerthelp.com For All NCERT solutions, CBSE sample papers, Question papers, Notes for Class 6 to 12



a) Draw the structure of B<sub>2</sub>H<sub>6</sub>.

5

- b) What happens when:
  - Boric acid is added to water.
  - ii) Aluminium is treated with dilute NaOH.
- c) Give suitable reason for the following:
  - [SiF<sub>6</sub>]<sup>2</sup> is known whereas [SiCl<sub>6</sub>]<sup>2</sup> not.
  - In group 14, the tendency for catenation decreases with increasing atomic number.

OR

- a) Complete the follwing chemical equations:
  - i)  $Fe_2O_3 + 3CO \xrightarrow{\Delta}$
  - ii) CaCO, +2HCl→
- b) Write a brief account on the following:
  - Diamond is covalent, yet it has high melting point.
  - ii) Atomic radius of gallium (135pm) is less than that of aluminium (143pm).
  - iii) Graphite is a good conductor of electricity but diamond is insulator.
- 30. a) Explain the following reactions with suitable examples:

5

- Wurtz reaction.
- ii) Friedal-Crafts alkylation reaction.
- An alkene 'A' on azonolysis gives mixture of ethanol and pentan-3 one.
   Write the structure and IUPAC name of 'A'.
- c) Give one chemical test to distinguish between ethene and ethyne.

OR

- a) Write suitable reason for the following:
  - C-C bond length in benzene ring is 139 pm which is in between C-C single bond 154 pm and C=C double bond 133 pm.
  - Trans-2-butene has higher melting point than cis-isomer.
- b) Give a chemical test to distinguish between but-1-yne and but-2-yne.
- c) How will you carry out the following conversions:
  - i) Ethene to benzene
  - ii) 1-bromopropane to 2-bromopropane.